

Radiance

PDO POST TREATMENT RECOMMENDATIONS:

- Avoid picking and manipulating the suture at the procedure site. This can cause more bruising, infection, the perforation of the suture through the surface of the skin or the disjunction of the suture to the skin and muscle.
- Avoid direct heat such as laser treatment such as IPL or laser hair removal for at least 2 weeks.
- Avoid excessive exercise/sport for one week following your procedure.
- For blood clot type bruises (hematomas) you may apply cold compresses gently to the area.
- Arnica or Vitamin K may be applied after procedure for 5-7 days for bruising and swelling if needed.
- There will be immediate visible results after treatment with suspensory threads, as well as the area feeling tighter. Over the first few days to weeks, the tight feeling will relax and the collagen will start building.
- Soreness always occurs after treatment. There may be a dull pain that typically goes away after a week or two. There will often be discomfort from chewing or any facial movement. Another kind of pain after PDO threads can be described as "twinges", "zingers", or "quick shocks", this is completely normal, and is caused by the threads stimulating a nerve ending. This kind of pain gradually resolves within several weeks after thread placement.
- Recovery is generally quick, with most patients returning to work the next day.
- Asymmetry or puckering of the skin may occur and typically resolves with time. If it does not resolve after a week, please contact us.
- You may take Tylenol for pain, especially at bedtime during the first few days.

When to call:

- If you experience increased redness, fever or drainage from the thread site.
- If a thread begins to poke through the skin.

Facial PDO Procedures:

- Keep head elevated on several pillows for the first 3 nights after procedure.
- Avoid any direct pressure to the face and area where the sutures were placed for one week after the procedure (for example, sleeping face down, excessive chewing, talking, yawning, smiling and other facial movements during the week after your procedure.
- You may wash your hair and face with very gentle upward and outward motions that lift the skin in the same directions as the sutures lift the skin.
- Patient can remove KT tape (if applied) after 24 hours. When removing KT tape, patients must remove carefully and in a bottom to top pulling direction. Moisten the area with a small amount of baby oil.
- Banding may appear, which is normal and is nothing to be concerned about. Applying warm water compresses and time will resolve any banding issues.

It is very important that you call us with concerns. Possible adverse reactions need to be reported to us immediately! A follow-up will be necessary if you are experiencing any type of skin reaction. During business hours call 608-240-0088. After business hours call Taylor Hill, RN @ 608-575-1750 or Anne Pientka, RN @ 608-513-8699.

Basecamp & Botox

BASECAMP FITNESS x *Radiance*

309 Junction Rd | 608.716.3700 | basecampfitness.com

Exclusive offer for Radiance clients

Get **1 FREE MONTH** of classes here:

